arXiv:1303.1056v1 [math.MG] 5 Mar 2013

On the synectic metric in the tangent bundle of a Riemannian manifold

Melek ARAS*

March 6, 2013

Abstract

The purpose of this paper is to investigate applications the covariant derivatives of the covector fields and killing vector fields with respect to the synectic lift ${}^{S}g = {}^{C}g + {}^{V}a$ in a the Riemannian manifold to its tangent bundle $T(M_n)$, where ${}^{C}g$-complete lift of the Riemannian metric, ${}^{V}a$-vertical lift of the symmetric tensör field of type $(0,2)$ in M_n .

Keywords: Tesor bundle ; Metric connection; Covector field; Levi-Civita connections; Killing vector field

1. Introduction

Suppose that there is given the following Riemannian metric

$$ {}^{S}\widetilde{g}_{CB}dx^{C}dx^{B} = a_{ji}dx^{j}dx^{i} + 2g_{ji}dx^{j}\delta y^{i} \tag{1}$$

in tangent bundle in $T(M_n)$ over a Riemannian manifold M_n with metric g, where a_{ji} are components of a symmetric tensor field of type $(0,2)$ in M_n and $\delta y^{h} = dy^{h} + \Gamma_{i}^{h}dx^{i}$, $\Gamma_{i}^{h} = y^{j}\ \Gamma_{ji}^{h}$ with respect to the induced coordinates (x^{h}, y^{h}) in $\pi^{-1}(U) \subset T(M_n)$. We call this metric the synectic metric. The synectic metric ${}^{S}g = {}^{C}g + {}^{V}a$ has respectively components[2] and its contravarient components

$$ {}^{S}g = \left({}^{S}\widetilde{g}_{CB}\right) = \begin{pmatrix} a_{ji} + \partial g_{ji} & g_{ji} \\ g_{ji} & 0 \end{pmatrix}, {}^{S}\widetilde{g}^{CB} = \begin{pmatrix} 0 & g^{ji} \\ g^{ji} & x^{\overline{s}}\partial_{s}g^{ji} - a^{ji}_{\cdot\cdot} \end{pmatrix} \tag{2}$$

where $\partial g_{ji} = x^{\overline{s}}\partial_{s}g_{ji}$ and $a^{ti}_{\cdot\cdot} = g^{jt}a_{js}g^{si}$.

Components of the Riemannian connection determined by the synectic metric ${}^{S}g$ are [1]

$$ \left\{ \begin{array}{l} {}^{S}\Gamma_{ji}^{k} = \Gamma_{ji}^{k}, \quad {}^{S}\Gamma_{\overline{j}i}^{\overline{k}} = \Gamma_{ji}^{k}, \quad {}^{S}\Gamma_{j\overline{i}}^{\overline{k}} = \Gamma_{ji}^{k}, \quad {}^{S}\Gamma_{\overline{j}\overline{i}}^{\overline{k}} = 0 \\ {}^{S}\Gamma_{j\overline{i}}^{k} = {}^{S}\Gamma_{\overline{j}i}^{k} = {}^{S}\Gamma_{\overline{j}\overline{i}}^{k} = 0, \quad {}^{S}\Gamma_{ji}^{\overline{k}} = x^{\overline{t}}\partial_{t}\Gamma_{ji}^{k} + H_{ji}^{k} \end{array} \right. \tag{3}$$

---

*Department of Mathematics,Faculty of Arts and Sciences, Giresun University, 28049, Turkey e-mail:melekaras25@hotmail.com;melek.aras@giresun.edu.tr with respect to the induced coordinates in $T\left(M_n\right)$, Γ_{ji}^{k} being Christoffel symbols constructed with g_{ji} . Where $H_{ji}^{k}=\frac{1}{2}g^{ks}\left(\nabla_j a_{si}+\nabla_i a_{js}-\nabla_s a_{ji}\right)$ is a tensor of type $(1,2)$ and $\nabla_s a_{ji}=\partial_s a_{ji}-\Gamma_{kj}^{l}a_{li}-\Gamma_{ki}^{l}a_{jl}$.

The metric connection $\widetilde{\nabla}$ of the synectic metric satisfies $\widetilde{\nabla}_C\ {}^S g_{BA}=0$ and has non-trivial torsion tensor $\overline{T}_{CB}^{A}$, which is skew-symmetric in the indices C and B .Then the metric connection $\widetilde{\nabla}$ of the synectic metric has components[1]

$$\begin{cases} \widetilde{\Gamma}_{ji}^{h}=\widetilde{\Gamma}_{\bar{j}i}^{\widetilde{h}}=\widetilde{\Gamma}_{j\bar{i}}^{\overline{h}}=\Gamma_{ji}^{h}, \\ \widetilde{\Gamma}_{ji}^{h}=\widetilde{\Gamma}_{\bar{j}i}^{\overline{h}}=\widetilde{\Gamma}_{j\bar{i}}^{\overline{h}}=\Gamma_{ji}^{h}, \\ \widetilde{\Gamma}_{ji}^{\overline{h}}=x^{t}\partial_t\Gamma_{ji}^{h}+H_{ji}^{h}-y^{k}R_{kjih} \end{cases} \tag{4}$$

with respect to the induced coordinates, Γ_{ji}^{k} being Christoffel symbols formed with g_{ji}, where $H_{ji}^{k}=\frac{1}{2}g^{ks}\left(\nabla_j a_{si}+\nabla_i a_{js}-\nabla_s a_{ji}\right)$.

Given a vector field $\widetilde{X}$ in $T\left(M_n\right)$, the $1-form$, $\widetilde{\omega}$ defined by $\widetilde{\omega}\left(\widetilde{Y}\right)=\widetilde{g}\left(\widetilde{X},\widetilde{Y}\right)$, $\widetilde{Y}$ being an arbitrary element of $T_0^1\left(M_n\right)$, is called the covector field associated with $\widetilde{X}$ and denoted by $\widetilde{X}^*$. If $\widetilde{X}$ has local components $\widetilde{X}^A$, then the associated covector field $\widetilde{X}^*$of $\widetilde{X}$ has local components $\widetilde{X}_C=\widetilde{g}_{CA}\widetilde{X}^A$.

Let ωbe a $1-form$ in M_n with components ω_i. Then the vertical , complete and horizontal lifts of ω to $T\left(M_n\right)$ have respectively components[4]

$$\left({}^V\omega_B\right)=\left(\omega_i,0\right),\qquad \left({}^C\omega_B\right)=\left(\partial\omega_i,\omega_i\right),\qquad \left({}^H\omega_B\right)=\left(-\Gamma_i^k\omega_k,\omega_i\right)$$

with respect to the induced coordinates in $T\left(M_n\right)$.

The associated covector fields of the vertical , complete and horizontal lifts to $T\left(M_n\right)$, with the synectic metric , of a vector field X with components X^h in M_n are respectively

$$\left(X_i,0\right),\quad \left(y^s\partial_s X_i+a_{ij}X^j,X_i\right),\quad \left(\Gamma_i^h X_h+a_{ij}X^j,X_i\right) \tag{5}$$

with respect to the induced coordinates, where $X_j=g_{ji}X^i$ are components of the covector field X^* associated with X.

A vector field $X\epsilon\Im_0^1\left(M_n\right)$ is said to be a Killing vector field of a Riemannian manifold with metric g, if $L_X g=0$[3]. In terms of components g_{ji} of g, X is a Killing vector field if and only if

$$L_X g=X^\alpha\nabla_\alpha g_{ji}+g_{\alpha i}\nabla_j X^\alpha+g_{j\alpha}\nabla_i X^\alpha=\nabla_j X_i+\nabla_i X_j=0,$$

X^α being components of X, where ∇ is the Riemannian connection of the metric g.

2. Main Results

We now take a vector field X in M_n with components X^h.Then, since the associated covector fields of the lifts of X have respectively components give by (5), we have by (3) and (5)

$$\left\{\begin{array}{l} {}^{S}\nabla^{V}_{B}X_{A}=\begin{pmatrix}\nabla_{j}X_{i} & 0\\ 0 & 0\end{pmatrix}, \\ {}^{S}\nabla^{C}_{B}X_{A}=\begin{pmatrix}\partial\left(\nabla_{j}X_{i}\right)+\nabla_{j}\left(a_{il}X^{l}\right)-H^{m}_{ji}X_{m} & \nabla_{j}X_{i}\\ \nabla_{j}X_{i} & 0\end{pmatrix}\end{array}\right. \tag{6}$$

and consequently

$$\left\{\begin{array}{l} {}^{S}\nabla^{V}_{B}X_{A}+^{S}\nabla^{V}_{A}X_{B}=\begin{pmatrix}\nabla_{j}X_{i}+\nabla_{i}X_{j} & 0\\ 0 & 0\end{pmatrix}, \\ {}^{S}\nabla^{C}_{B}X_{A}+^{S}\nabla^{C}_{A}X_{B}=\begin{pmatrix}{}^{S}\nabla^{C}_{j}X_{i}+^{S}\nabla^{C}_{i}X_{j} & {}^{S}\nabla^{C}_{j}X_{\bar{i}}+^{S}\nabla^{C}_{\bar{i}}X_{j}\\ {}^{S}\nabla^{C}_{\bar{j}}X_{i}+^{S}\nabla^{C}_{i}X_{\bar{j}} & {}^{S}\nabla^{C}_{\bar{j}}X_{\bar{i}}+^{S}\nabla^{C}_{\bar{i}}X_{\bar{j}}\end{pmatrix} \\ {}^{S}\nabla^{C}_{j}X_{i}+^{S}\nabla^{C}_{i}X_{j}=\partial\left(\nabla_{j}X_{i}+\nabla_{i}X_{j}\right) \\ \qquad\qquad+\nabla_{j}\left(a_{il}X^{l}\right)+\nabla_{i}\left(a_{jl}X^{l}\right)-H^{m}_{ji}X_{m}-H^{m}_{ij}X_{m} \\ \quad{}^{S}\nabla^{C}_{j}X_{\bar{i}}+^{S}\nabla^{C}_{\bar{i}}X_{j}=\nabla_{j}X_{i}+\nabla_{i}X_{j} \\ \quad{}^{S}\nabla^{C}_{\bar{j}}X_{i}+^{S}\nabla^{C}_{i}X_{\bar{j}}=\nabla_{j}X_{i}+\nabla_{i}X^{S}_{j} \\ \qquad\nabla^{C}_{\bar{j}}X_{\bar{i}}+^{S}\nabla^{C}_{\bar{i}}X_{\bar{j}}=0 \\ \end{array}\right. \tag{7}$$

with respect to the induced coordinates ,where $X_{j}=g_{jk}X^{k}$. From (7) we have

Theorem 1 *Necessery and sufficient conditions in order that* (a) *the vertical,* (b) *complete lifts to* $T\left(M_{n}\right)$*, with the synectic metric, of a vector field* X *in* M_{n} *be a Killing vector field in* $T\left(M_{n}\right)$ *are that, respectively,* (a) X *is a Killing vector field in* M_{n}*and* (b) X *is Killing vector field with vanishing covariant derivative in* M_{n} *and the covariant derivative of symmetric tensor field* a *of type* $(0,2)$ *vanishes.*

We also have by (2) and (6)

$$\left\{\begin{array}{l} {}^{S}\nabla^{V}_{B}X_{A}-^{S}\nabla^{V}_{A}X_{B}=\begin{pmatrix}\nabla_{j}X_{i}-\nabla_{i}X_{j} & 0\\ 0 & 0\end{pmatrix}, \\ {}^{S}\nabla^{C}_{B}X_{A}-^{S}\nabla^{C}_{A}X_{B}=\begin{pmatrix}{}^{S}\nabla^{C}_{j}X_{i}-^{S}\nabla^{C}_{i}X_{j} & {}^{S}\nabla^{C}_{j}X_{\bar{i}}-^{S}\nabla^{C}_{\bar{i}}X_{j}\\ {}^{S}\nabla^{C}_{\bar{j}}X_{i}-^{S}\nabla^{C}_{i}X_{\bar{j}} & {}^{S}\nabla^{C}_{\bar{j}}X_{\bar{i}}-^{S}\nabla^{C}_{\bar{i}}X_{\bar{j}}\end{pmatrix} \\ {}^{S}\nabla^{C}_{j}X_{i}-^{S}\nabla^{C}_{i}X_{j}=\partial\left(\nabla_{j}X_{i}-\nabla_{i}X_{j}\right) \\ \qquad\qquad+\nabla_{j}\left(a_{il}X^{l}\right)-\nabla_{i}\left(a_{jl}X^{l}\right)-H^{m}_{ji}X_{m}+H^{m}_{ij}X_{m} \\ \quad{}^{S}\nabla^{C}_{j}X_{\bar{i}}+^{S}\nabla^{C}_{\bar{i}}X_{j}=\nabla_{j}X_{i}-\nabla_{i}X_{j} \\ \quad{}^{S}\nabla^{C}_{\bar{j}}X_{i}+^{S}\nabla^{C}_{i}X_{\bar{j}}=\nabla_{j}X_{i}-\nabla_{i}X^{S}_{j} \\ \qquad\nabla^{C}_{\bar{j}}X_{\bar{i}}-^{S}\nabla^{C}_{\bar{i}}X_{\bar{j}}=0 \\ \end{array}\right. \tag{8}$$

with respect to the induced coordinates ,where ${}^{S}g^{BA}\,{}^{S}\nabla^{V}_{B}X_{A}=0$ and ${}^{S}g^{BA}\,{}^{S}\nabla^{C}_{B}X_{A}=2g^{ji}\nabla_{j}X_{i}$. Thus we have, from (8), respectively,

Theorem 2 *The vertical lift of a vector field in* M_n *to* $T(M_n)$ *with the synectic metric* ${}^S g$ *is harmonic if and only if the vector field in* M_n *is closed.*

Theorem 3 *The complete lift of a vector field in* M_n *to* $T(M_n)$ *with the synectic metric* ${}^S g$ *is harmonic if and only if the vector field in* M_n *is harmonic and the covariant derivative of symmetric tensor field a of type* $(0,2)$ *vanishes.*

We consider a vector field $X \epsilon \Im_0^1(M_n)$. Then its vertical , complete and horizontal lifts have components of the form

$$ {}^V X = \begin{pmatrix} 0 \\ X^h \end{pmatrix}, \quad {}^C X = \begin{pmatrix} X^h \\ \partial X^h \end{pmatrix}, \quad {}^H X = \begin{pmatrix} X^h \\ -\Gamma_i^h X^i \end{pmatrix} \tag{9} $$

with respect to the induced coordinates in $T(M_n)$, where $\Gamma_i^h X^i = y^s \ \Gamma_{si}^h X^i$.

Let X be a vector field in M_n with local components X^k. Then, from (9) and (4), we see that, the covariant derivatives of the vertical, complete and horizontal lifts of $X \epsilon \Im_0^1(M_n)$with the metric connection $\widetilde{\nabla}$ have respectively components

$$ \left\{ \begin{array}{c} \widetilde{\nabla}_B \ {}^V X^A = \begin{pmatrix} 0 & 0 \\ \nabla_j X^h & 0 \end{pmatrix}, \\ \widetilde{\nabla}_B \ {}^C X^A = \begin{pmatrix} \nabla_j X^h & 0 \\ \partial\left(\nabla_j X^h\right) + H_{jm}^h X^m - y^k R_{kjm}^h X^m & \nabla_j X^h \end{pmatrix}, \\ \widetilde{\nabla}_B \ {}^H X^A = \begin{pmatrix} \nabla_j X^h & 0 \\ -\Gamma_i^h \left(\nabla_j X^i\right) + H_{jm}^h X^m & 0 \end{pmatrix} \end{array} \right. \tag{10} $$

with respect to the induced coordinates $T(M_n)$.

Remark 4 $\widetilde{\nabla} = \overline{\nabla} +^V H$, *where* $\overline{\nabla}$ *is the metric connection with the metric* ${}^C g$*[1]*

Remark 5 *The metric connection* $\overline{\nabla}$ *coincides with the horizontal lift* ${}^H\nabla$ *of Levi Civita connection* ∇*of* g *in* M_n*[4] . Thus we have*

Proposition 6 *Necessery and sufficient conditions in order that* (a) *the vertical,* (b) *complete and horizontal lifts of a vector field in* M_n *to* $T(M_n)$ *with the metric connection* $\widetilde{\nabla}$ *be parallel in* $T(M_n)$ *are that, respectively,* (a) *the vector field given in* M_n *is parallel* (b) *the vector field given in* M_n *is parallel and the covariant derivative of symmetric tensor field a of type* $(0,2)$ *vanishes.*

Since $\nabla_j X^h = t\delta_j^h$ with constant t implies $R_{kji}^h X^i = 0$, we have also

Proposition 7 *The complete lift of a vector in* M_n *to* $T(M_n)$ *with the metric connection* $\widetilde{\nabla}$*is concurrent if and only if the vector field given in* M_n *is concurrent and the covariant derivative of symmetric tensor field a of type* $(0,2)$ *vanishes.*

A vector field $X \epsilon \Im_0^1 \left(M_n \right)$ is said to be an infinitesimal isometry or a Killing vector field of a Riemannian manifold with metric g, if $L_X g = 0$. In terms of components g_{ji} of g, X is an infinitesimal isometry if and only if

$$X^\gamma \partial_\gamma g_{\alpha\beta} + g_{\alpha\gamma} \partial_\beta X^\gamma + g_{\gamma\beta} \partial_\alpha X^\gamma = 0, \tag{11}$$

X^α being components of X[3], where the indices α, β and γ run over the range $\{1, 2, ..., m\}$.

Let there be given in M_n a Riemannian metric g with components g_{ji}.Let $\widetilde{X}$ be a vector field with components $\begin{pmatrix} \widetilde{X}^k \\ \widetilde{X}^{\overline{k}} \end{pmatrix}$ with respect to the induced coordinates in $T \left(M_n \right)$.

with respect to the induced coordinates in $T \left(M_n \right)$. Then, taking account of (2), we see by virtue of (11) that $\widetilde{X}$ is an infinitesimal isometry n $T \left(M_n \right)$ with metric ${}^S g$ if and only if

$$\left\{ \begin{array}{l} \left(\widetilde{X}^h \partial_h \partial g_{ji} + \widetilde{X}^{\overline{h}} \partial_h g_{ji} \right) + \left(\partial g_{jh} \partial_i \widetilde{X}^h + g_{jh} \partial_i \widetilde{X}^{\overline{h}} \right) \\ + \left(\partial g_{hi} \partial_j \widetilde{X}^h + g_{hi} \partial_j \widetilde{X}^{\overline{h}} \right) + \left(\widetilde{X}^h \partial_h a_{ji} + a_{jh} \partial_i \widetilde{X}^h + a_{hi} \partial_j \widetilde{X}^h \right) = 0 \end{array} \right. \tag{12}$$

$$\widetilde{X}^h \partial_h g_{ji} + g_{jh} \partial_i \widetilde{X}^h + \left(\partial g_{hi} \partial_{\overline{j}} \widetilde{X}^h + g_{hi} \partial_{\overline{j}} \widetilde{X}^{\overline{h}} \right) + a_{hi} \partial_{\overline{j}} \widetilde{X}^h = 0 \tag{13}$$

$$\widetilde{X}^h \partial_h g_{ji} + g_{hi} \partial_j \widetilde{X}^h + \left(\partial g_{jh} \partial_{\overline{i}} \widetilde{X}^h + g_{jh} \partial_{\overline{i}} \widetilde{X}^{\overline{h}} \right) + a_{jh} \partial_{\overline{i}} \widetilde{X}^h = 0 \tag{14}$$

$$g_{jh} \partial_{\overline{i}} \widetilde{X}^h + g_{hi} \partial_{\overline{j}} \widetilde{X}^h = 0. \tag{15}$$

We shall now prove

Lemma 8 *Let C be an element of $\Im_1^1 \left(M_n \right)$. Then $L_{\iota C} \ {}^S g = 0$ holds if and only if $C = 0$.*

Proof. *Denote by C_i^k the local components of C. Then ιC has components $\begin{pmatrix} 0 \\ y^i C_i^k \end{pmatrix}$ with respect to the induced coordinates in $T \left(M_n \right)$.Thus, substituting $\widetilde{X}^k = 0$ and $\widetilde{X}^{\overline{k}} = y^i C_i^k$ in (13) , we have $g_{ki} C_j^k = 0$, which implies $C_j^k = 0$, i.e., $C = 0$.* ■

References

[1] Aras, M. The Metric Connection with Respect to the Synectic Metric. Hacet J. Math. Stat, 2012, 41(2), 169-173.

[2] Vishnevskii V V, Shirokov A and Shurygin V V, Spaces over algebras (Kazan University Press) (1985)

[3] Yano, K. The Theory of Lie Derivatives and Applications, Elsevier, Amsterdam, 1957.

[4] Yano, K. and Ishıhara, S. Tangent and Cotangent Bundles, Marcel Dekker Inc. New York, 1973.